

Quality of commercial soybean grain and post-harvest economic losses in farming in West Pará, Brazil

Juliana Amaral Vinhote¹, Gabriel Coradini¹, Éder Bruno Rebelo da Silva², Welligton Conceição da Silva², Antônio Vinicius Correa Barbosa², Maria Roseane Pereira dos Santos³, Gilbson Santos Soares^{1,4}

¹Universidade Luterana do Brasil – ULBRA. ²Universidade Federal Rural da Amazônia - UFRA, PA. ³Universidade Federal do Oeste do Pará – UFOPA, PA. ⁴Instituto Federal do Pará – IFPA, PA. E-mail: julianavinhote@rede.ulbra.br

Abstract

The soybean production chain is one of the main agribusiness products. In this context, the grain market is constantly expanding. Thus, it is important to carry out the characterization of soybeans and define their qualities. The objective of this study was to evaluate the quality of commercial soybean grains in post-harvest, as well as the economic losses in a crop in the municipality of Mojuí dos Campos, Pará. Two types of cultivars at 4280 (IPRO) and 8644 (IPRO) were used, harvested on different days and plots. Data collection was carried out by monitoring the post-harvest process. The information regarding the grains was obtained through the classification report provided by the company accredited to classify the grains. The methods used were carried out in accordance with the Technical Regulation of Soybeans of Normative Instruction No. 11, of May 15, 2007, of the Ministry of Agriculture, Pecuária e Abastecimento (MAPA). Data were analyzed by PROC TTEST in SAS version 19.0 at a significance level of 5%. The results showed that, in the period of harvesting of cultivar 8644(IPRO), there was a higher rate of rainfall compared to the days of harvesting of cultivar 4280(IPRO). The total of discounted grains was 15.101t for the cultivar 4280 (IPRO). The wet grain is the only parameter responsible for the discount in both cultivars. The total of discounted grains referring to cultivar 8644 (IPRO) was 30.218t. The wet grains for the cultivar had the highest classification rate on the fourth day of harvest, with a total of 23.20%, being related to the decrease in the percentage of broken grains (1.60%) and a higher index of burnt/damaged grains. It is concluded that the cultivar 4280 (IPRO) presented a lower grain discount compared to the cultivar 8644 (IPRO). Consequently, cultivar 8644 presented greater economic damage to the property.

Keywords: quantitative classification of grains; moisture; *Glycine max*; qualitative classification of grains.

Qualidade de grãos comerciais de soja e perdas econômicas na pós-colheita em lavoura no Oeste do Pará, Brasil

RESUMO

A cadeia produtiva da soja é um dos principais produtos do agronegócio. Nesse contexto, o mercado de grãos está em constante expansão. Assim, faz-se importante realizar a caracterização dos grãos de soja e definir as suas qualidades. O objetivo neste estudo foi avaliar a qualidade de grãos comerciais de soja na pós-colheita, bem como as perdas econômicas em uma lavoura no município de Mojuí dos Campos, Pará. Foram utilizados dois tipos de cultivares a 4280^(IPRO) e 8644^(IPRO), colhidas em diferentes dias e parcelas. A coleta dos dados foi realizada por meio do acompanhamento do processo de pós-colheita. As informações referentes aos grãos foram obtidas por meio do Laudo de classificação fornecido pela empresa credenciada a classificar os grãos. Os métodos empregados foram realizados conforme o Regulamento Técnico da Soja da Instrução Normativa Nº 11, de 15 de maio de 2007, do Ministério da Agricultura, Pecuária e Abastecimento (MAPA). Os dados foram analisados pelo PROC TTEST no SAS versão 19.0 a um nível de significância de 5%. Os resultados mostraram que, no período de colheita da cultivar 8644^(IPRO), houve um maior índice de precipitação pluvial em comparação com os dias de colheita da cultivar 4280^(IPRO). O total de grãos descontados foi de 15,101t para a cultivar 4280^(IPRO). O grão úmido é o único parâmetro responsável

peelo desconto em ambas as cultivares. O total de grãos descontados referentes à cultivar 8644^(IPRO) foi de 30,218t. Os grãos úmidos para a cultivar obtiveram maior taxa de classificação no quarto dia da colheita com total de 23,20%, tendo relação com a diminuição da porcentagem dos grãos quebrados (1,60%) e um maior índice de grãos ardidados/avariados. Conclui-se que a cultivar 4280^(IPRO), apresentou menor desconto de grãos em comparação com a cultivar 8644^(IPRO). Por consequência a cultivar 8644 apresentou maior dano econômico à propriedade.

Palavras-chave: classificação quantitativa de grãos; umidade; *Glycine max*; classificação qualitativa de grãos.

Introduction

Soybean (*Glycine max*) belongs to the Fabaceae (legume) family, from the Asian continent, in the region of ancient China, its grain has a high protein content and a large amount of essential amino acids for the functioning of the human body (POLLNOW *et al.*, 2020). In Brazil, soybean crops reach 37.03 million planted hectares, representing a production of 126 million tons in the 2019 harvest (CONAB, 2020).

In national and global agribusiness, soy is one of the crops of great economic importance. This fact favors the development and structuring of the international market, the consolidation of the grain as a source of vegetable protein, and the emergence of new technologies that enabled the expansion of exploration in different regions of the world (HIRAKURI; LAZZAROTTO, 2014; BUSS *et al.*, 2019).

Soybeans suffer several interferences in their quality, which can occur in the field, during harvesting and drying, in processing and storage, transport and sowing (FRANÇA NETO *et al.*, 2016). Factors that can intensify due to temperature extremes during maturation, fluctuations in relative air humidity, drought, nutritional and water deficit, occurrence of pests and diseases, in addition to inadequate procedures during harvesting, drying and storage, in addition, the excess of moisture causes deterioration before harvesting, which can result in high rates of mechanical damage (VINHOTE *et al.*, 2021).

The quality of soy beans is the main parameter for their commercialization and processing, because when they occur inappropriately, they affect the price of the product, a fact that shows that Brazilian agriculture still does not fully control losses during the post-harvest, as they suffer interference of external factors, being physical, such as temperature, humidity; chemical, such as

oxygen supply, and biological, such as bacteria, fungi, insects and rodents (SAATH *et al.*, 2017).

Soybean classification takes place in accordance with Normative Instruction No. 11, of May 15, 2007, of the Ministry of Agriculture, Pecuária e Abastecimento (BRASIL, 2007a; 2007b), which standardizes defects, rules and limits of soy beans for the commercialization. The main patterned defects are: burnt grains, hot grains, moldy grains, fermented grains, sprouted grains, immature grains, damaged grains, empty grains. Being classified as greenish, the grains or pieces of grain with complete physiological development. Damaged grains represent the sum of burnt, hot, moldy, fermented, germinated, immature, damaged by pests and empty grains, with a tolerance of 8%. Moist grains, those with a variable proportion of water inside, have a tolerance of 14% (BRASIL, 2007a).

Thus, soybeans for commercialization must present normative standards, seeking quality models for the grains and adequate processes during their processing, which starts at harvest and extends throughout the entire cycle, until its processing, aiming to reduce losses and reduce the injuries on the grain. In this context, this study is justified by the need to identify, according to official standards, the characteristics of soybeans subject to commercialization, observing whether market requirements are met with aptitude and homogeneous quality, spotting the main injuries present in Soy.

In addition, in the post-harvest, the classification of grains is essential, as it will present more consistent responses, which may be physical, physiological or mechanical, encouraging the scientific community to use this tool and aiming to address in a timely manner the origin of the damage caused by the production of soybeans in Brazil (VINHOTE *et al.*, 2021).

The hypothesis of this study was that there are differences in the quality of soybeans


harvested on different days when comparing two types of cultivars 4280(IPRO) and 8644(IPRO). Based on this information, the aim of this study was to evaluate the quality of commercial soybean grains in post-harvest, as well as the economic losses in a crop in the municipality of Mojuí dos Campos, Pará.

Material and Methods

The study was carried out in a soybean plantation, located in the municipality of Mojuí dos Campos (02° 10' 17" S and 56° 44' 42" W), in

the north of Pará. With a territorial area equivalent to 4,988,236 km², the soybean crop has a total area of 124.63 ha, (59.46 - 4280) (65.17 - 8644) which was divided into plots according to the corresponding cultivar and collection day, which is shown in Figure 1.

Figure 1. Location map of harvesting areas, Mojuí dos Campos, Pará.



In this study, two types of cultivars were used, the 4280(IPRO) and the 8644(IPRO). The first has greater tolerance to rain at the time of harvest.

Both cultivars were planted and harvested on the same type of soil. The soil texture was characterized by the combination of clay, silt and sand. Among those, clay was the one that presented the greatest quantity. According to the soil analysis report carried out on the property, the soil can be classified as very clayey.

And so, the necessary corrections, fertilization and cultural treatments were made to receive sowing, according to technical

recommendations, thus adopting the spacing between rows of 0.45 m, density of 10 plants/m⁻¹ for the cultivars.


Data collection was carried out for three days by monitoring the harvest and post-harvest process until the transfer to the accredited company responsible for grain classification.

The harvest of both cultivars took place on different days and plots (Figures 2 and 3), being fully mechanized, and the harvester displacement speed was 6 km/h⁻¹.

Figure 2. Cultivar harvest area, photographed by Drone (Brand Xiaomi, model FIMI X8 SE).



Figura 3. Location map of harvesting areas for cultivar 4280(IPRO) and 8644(IPRO), Mojuí dos Campos, Pará.



Information regarding rainfall during harvest of soybean cultivars was also collected. For this, a rain gauge was used.

Information related to the grains were obtained through a classification report provided by the company authorized to classify the soy grains destined for commercialization, based on their intrinsic and extrinsic qualities.


The methods used were carried out in accordance with the Technical Regulation of Soybean Normative Instruction No. 11, of May

15, 2007, of the Ministry of Agriculture, Pecuária e Abastecimento (MAPA), referring to sampling and the process of obtaining these, standardized with the values prefixed.

The parameters evaluated are qualitative, which are characterized according to the defects, and quantitative, which are related to the levels of tolerance of the defects found, which are displayed in percentages.

Classification of soy beans was carried out in compliance with the current Normative

Instruction following the chronological order as shown in the flowchart below:



To calculate the percentage of damaged grains, the formula was adopted:

$$\% = \frac{\text{damaged grain weight (g)}}{\text{sample weight}} \times 100$$

The total value of damaged groups represented by burnt (Q), scorched (A), moldy (M), fermented (F), germinated (G), damaged (D), immature (I) and empty (C) will be determined by middle of the formula:

$$\sum (Q + A + M + F + G + D + I + C) = X$$

To determine the percentage of foreign matter and impurities - MEI, the formula will be used:

$$\% = \frac{MEI \text{ weight (g)}}{\text{sample weight}} \times 100$$

To estimate the economic loss caused by the discount of wet grains, the total weight of the discounted grains was used, in relation to the value, in reais, practiced during the month of May 2021, equivalent at the time of the research to R\$ 160.00/sack.

To calculate the economic loss, the following formula was used:

1ª) Discounted total weight / sack weight (60 kg)


2ª) Sack value vs. total discounted sacks

After collection, data were tabulated and later organized in Microsoft Office Excel® spreadsheets (2016). These were analyzed statistically by PROC TTEST in SAS version 19.0 at a significance level of 5%.

Results and Discussion

Regarding rainfall, it was found that during the harvest period of cultivar 8644(IPRO), there was a higher rate of rainfall compared to the days of harvest of cultivar 4280(IPRO), as seen in Figure 4.

Figure 4. Rainfall data in the four days of collection for each cultivar.



The total of discarded grains was 15.101t, which corresponds to an index of 7.42% grain discount due to damage to the soybeans for cultivar 4280 (IPRO), with the wet grain being the only parameter responsible for the discount. Table 1 shows the classification of grains.

The total of discarded grains referring to cultivar 8644(IPRO) was 30.218t, which

corresponds to an 11.24% grain discount rate due to general damage to soybean grains.

Table 1 shows the classification of grains. As also observed in the cultivar (4280IPRO), the wet grain was the only parameter responsible for the grain discount.

Table 1. Classification of grains, Mojuí dos Campos, Pará.

Data in % (Cultivar 4280 ^(IPRO))					
Day	Moisture	Impurities	Hot/Damaged	Broken	Greenish
1	17,16	0,30	2,46	9,16	0
2	18,75	0,65	2,40	8,10	0
3	19,70	0,60	2,85	4,95	0
4	23,20	0,30	4,70	1,60	0
Data in % (Cultivar 8644 ^(IPRO))					
1	21,55	0,75	6,30	6,00	0
2	22,90	0,30	2,55	2,80	0
3	18,43	0,63	3,13	4,40	0
4	19,15	0,70	4,10	4,35	0

Thus, wet grains had a higher classification index on the fourth day of harvest,

with a total of 23.20%, related to the decrease in the percentage of broken grains (1.60%) and a

higher index of burnt/damaged grains. In accordance with the data from the present study, Lacerda *et al.* (2003) evidenced the dependence between grain moisture and rainfall, thus requiring the chemical desiccation process in order to dehydrate the grains, since the high moisture content in the studied plots would not be ideal for the moment of harvest. Therefore, it is noted that the water content is extremely important, as described by Botelho *et al.*, 2016, Santos *et al.*, 2016 e Khan *et al.*, 2017.

When there was a lower moisture rate (17.16%) there was an increase in the broken grain index (9.16%). Corroborating this study, Marcondes *et al.* (2005) verified the relationship between soybean grain moisture and low rainfall, observing grains with a lower percentage of moisture. It is noteworthy that the presence of broken grains is more evident when the grain has less moisture, making it more easily breakable (CARRARO *et al.*, 1985). Grains harvested with humidity above 18% may show damage, especially during threshing. On the other hand, grains with a moisture content below the recommended become more sensitive to breakage, thus presenting higher levels of broken grains (FRANÇA NETO *et al.*, 2007).

In cultivar 8644 (IPRO) it was also possible to observe a high moisture content (22.90%) on the second day, being related to the decrease in the broken indexes (2.80%). Corroborating the data described in this study, Borges *et al.* (2006) described that the delay in harvesting on grain yield caused significant losses, which were mainly influenced by the rainfall and genetic characteristics of the cultivars. Furthermore, late harvesting can lead to the deterioration of the physiological quality of the grains (DINIZ *et al.*, 2013; XAVIER *et al.*, 2015). This fact was evidenced in the harvest period of cultivar 8644 (IPRO), since it was delayed due to mechanical failures of the harvester, followed by high rainfall, thus delaying the ideal time for grain harvest.

The high losses observed during harvesting in this study may have occurred mainly due to the high frequency of rainfall (TSUKAHARA *et al.*, 2016). This fact was evidenced in the harvest of the 8644 (IPRO) culture, in which the rainfall index was more intense (Figure 4). Sales (2017) reinforces that the high frequency of rainfall, associated with fluctuations in relative humidity and temperature fluctuations, lead to losses in the physical,

physiological and sanitary quality of the grains. Still in this context, Borém (1999) states that soybean is characterized as an annual crop that presents variability of morphological characteristics, being directly influenced by climatic variables.

The classification showed indexes of burnt/damaged grains and the presence of impurities, all within the accepted range. During classification, burned grains are considered as one of the most important defects, as the fermentation process results in the browning of the grain, reducing its commercial quality (TEIXEIRA, 2001; BRASIL, 2007b). According to Souza (2012), when storing grains with a high percentage of burnt, that is, above acceptable limits, there are not only economic losses, as burnt grains have mycotoxins that in small quantities can affect human and animal health.

In the present study, the presence of impurities was evidenced in the volumes of analyzed samples. According to Lourenço (2019), less than 1% of impurities is associated with a good adjustment of the machine at the time of harvest, indexes greater than 1% make it more difficult to process the grains, making it necessary to go through more sieves. Botelho *et al.* (2019) describe that impurities are classified by the attribute of the physical quality of the grain, which refers to the quality of each batch by the grain and inert material index, number of grains of other species and harmful grains tolerated and prohibited.

It was also evidenced that both cultivars did not show greenish grain rate in the classification. Thus, França Neto *et al.* (2012) state that greenish beans are soybeans that have not reached their physiological development, which are subject to biotic or abiotic stresses, which, due to the application of products, end up presenting premature death or forced maturation, producing greenish beans with low physiological quality. The presence of greenish grains occurs due to the occurrence of inadequate temperatures, which accelerate the translocation of photoassimilates with lower rates of photosynthesis, which block the degradation of chlorophyll, resulting in greenish and poor quality grains (MARCOS FILHO, 2005).

In addition, greenish beans present high deterioration rates, which can result in reduced germination and vigor of soybean lots (FRANÇA NETO *et al.*, 2016).

There was the presence of damaged grains, specifically hot ones. In this context, damaged grains represent the sum of the grains that suffered damages during the harvesting process, comprising the hot, moldy, fermented, damaged, immature, hollow, germinated and burnt grains, with a tolerance of 8% in the commercialization of grains.

In this study, the economic losses identified by the grain discount correspond, in current values, to R\$ 120,850.66. Of these, the greatest economic loss corresponded to cultivar 8644(IPRO) (Table 2).

Table 2. Grains and economic losses in reais, Mojuí dos Campos, Pará, Brazil.

Cultivar	Sack weight (kg)	Sack value (R\$)	Economic loss (R\$)
4280 ^(IPRO)	60	160,00	40.269,33
8644 ^(IPRO)	60	160,00	80.581,33
Total	-	-	120.850,66

When performing the statistics of the data regarding the classification of grains for the two cultivars under study, it was observed that there was no significant difference ($p > 0.05$) between the studied variables, that is, this result indicates that both cultivars have similarity in

classification of grains, even with the difference existing in the days of harvest regarding rainfall (Table 3).

Table 3. Grain classification statistical data for cultivar 4280(IPRO) and 8644(IPRO).

Variables	P value
Moisture	0,6380
Impurity	0,2070
Hot/Damaged	0,0860
Broken	0,4236

Note = p value greater than 0.05 has no statistical difference.

Conclusion

Based on this information, it was possible to identify that cultivar 4280(IPRO) had a lower grain discount compared to cultivar 8644(IPRO). Consequently, cultivar 8644 presented greater economic damage to the property. Regarding impurities, it was noted that the variable moisture exceeded the acceptable limits in both cultivars, it was also found that the days with higher rainfall caused influence on moisture, resulting in a reduction in the percentage of broken grains.

There were no greenish grains, however, the classification showed indexes of hot/damaged grains and the presence of impurities, all within the accepted range, and moisture was the only factor that contributed to the discount of soybeans, causing considerable economic losses for the producer.

As an intervention measure, the use of grain dryers on the property is recommended, in order to prevent the soy beans from retaining

high concentrations of water, which, as a consequence, may cause them to be discounted.

References

- BORÉM, A. Escape gênico. **Biotecnologia, Ciência e Desenvolvimento (Encarte Especial)**, p.101-107, 1999.
- BORGES, I. O.; MACIEL, A. J. S.; MILAN, M. Programa computacional para o dimensionamento de colhedoras considerando a pontualidade na colheita de soja. **Engenharia Agrícola**, v.26, p.131-141, 2006. <https://doi.org/10.1590/S0100-69162006000100015>
- BRASIL. Ministério da Agricultura, Pecuária e Abastecimento. Instrução normativa n. 11, de 15 de maio de 2007. Estabelece o Regulamento Técnico da Soja, definindo o seu padrão oficial de classificação, com os requisitos de identidade e qualidade intrínseca e extrínseca, a amostragem

e a marcação ou rotulagem. **Diário Oficial da República Federativa do Brasil**, Brasília, n.93, p.13-15, 16 maio 2007a. Seção 1.

BRASIL. Ministério da Agricultura, Pecuária e Abastecimento. Instrução normativa n. 37, de 27 de julho de 2007. Altera o inciso IV, do art. 2º, do Capítulo I, do anexo da Instrução Normativa n. 11, de 15 de maio de 2007, que passa a vigorar com alterações, dando-se nova redação às alíneas “b” e “g” e acrescentando-se a alínea “h”. **Diário Oficial da República Federativa do Brasil**, Brasília, n. 145, p. 9, 30 jul. 2007b. Seção 1.

BUSS, R. N.; MENDANHA, J. F.; SILVA, D. M.; SIQUEIRA, M. G. Infraestrutura logística de transporte e armazenagem da soja no estado do Maranhão – Brasil. **Brazilian Journal of Development**, v.5, n.12, p.31564-31580, 2019. <https://doi.org/10.34117/bjdv5n12-247>

CARRARO, I. M.; BECO, A.; ROCHA, A. Efeito do retardamento da colheita sobre a qualidade de grãos de soja em Palotina, PR. **Revista Brasileira de Grãos**, v.7, p.123-132, 1995. <https://doi.org/10.17801/0101-3122/rbs.v7n3p123-132>

CONAB. **Acompanhamento da safra brasileira: grãos - Safra 2019**. Brasília, 2020. Disponível em: <https://www.conab.gov.br/conteudos.php?a=1253&t=.pdf>. Acesso em: 27 Jul. 2021.

DINIZ, F. O.; REIS, M. S.; DIAS, L. A. S.; ARAÚJO, E. F.; SEDIYAMA, T.; SEDIYAMA, C. A. Physiological quality of soybean seeds of cultivars submitted to harvesting delay and its association with seedling emergence in the field. **Journal of Seed Science**, v.35, p.147-152, 2013. <https://doi.org/10.1590/S2317-15372013000200002>

FRANÇA NETO, J. B.; KRZYZANOWSKI, F. C.; HENNING, A. A.; PÁDUA, G. P.; LORINI, I.; HENNING, F. A. **Tecnologia da produção de grão de soja de alta qualidade**. Londrina: Embrapa Soja, 2016. 84 p (Circular técnico. Documento; 380).

FRANÇA NETO, J. B.; KRZYZANOWSKI, F.C.; PÁDUA, P.; COSTA, N.P.; HENNING, A. **Tecnologia da produção de grão de soja de alta qualidade – Série Grãos**. Londrina: Embrapa Soja, 2007. 12p. (Embrapa Soja. Circular técnica, 40).

FRANÇA NETO, J. B.; PÁDUA, G. P.; KRZYZANOWSKI, F. C.; CARVALHO, M. L. M.; HENNING, A. A.; LORINI, I. **Grão esverdeada de soja: causas e efeitos sobre o desempenho fisiológico**. Londrina: Embrapa, 2012. (Série Grãos; n.91, 16 p.)

HIRAKURI, M. H.; LAZZAROTTO, J. J. O agronegócio da soja nos contextos mundial e brasileiro. **Documentos**, v. 349, p. 1-70, 2014.

LACERDA, A. L. S.; LAZARINI, E.; SÁ, M. E.; VALÉRIO FILHO, V. V. Aplicação de dessecantes na cultura de soja: teor de umidade nas grãos e biomassa nas plantas. **Planta Daninha**, v.21, p.427-434, 2003. <https://doi.org/10.1590/S0100-83582003000300011>

LOURENÇO, W. C. **Estudo de caso: Caracterização de cargas de grãos de soja da empresa Chapada Grãos, na safra 2018/2019**. 2019. Monografia (Graduação) - Universidade Federal de Santa Catarina, Campus Curitibanos Agronomia, 2019.

MARCONDES, M. C.; MIGLIORANZA, E.; FONSECA, I. C. B. Danos mecânicos e qualidade fisiológica de grão de soja colhida pelo sistema convencional e axial. **Revista Brasileira de Grãos**, v.27, p.125-129, 2005. <https://doi.org/10.1590/S0101-31222005000200018>

MARCOS FILHO, J. **Fisiologia de grãos de plantas cultivadas**. Piracicaba: Fealq, 2005. 495 p.

POLLNOW, H. E.; PIMENTEL, J. R.; POLLNOW, C.; PETER, M.; MEDEIROS, L. B.; PETER, M.; AUMONDE, T. Z.; PEDÓ, T. Manejo da adubação de base em soja no Noroeste do Rio Grande do Sul. **Brazilian Journal of Development**, v.6, n.6, p.38913-38923, 2020. <https://doi.org/10.34117/bjdv6n6-435>

SAATH, R.; SILVA TAVEIRA, J. H.; TERCENIANO, R. M.; THIAGO EVARISTO, T.; DEL ROSSO, B. C. Desempenho fisiológico e qualidade físico-química de grãos de soja secados sob diferentes temperaturas. **Colloquium Agrariae**, v.13, n.2, p.19-33, 2017. <https://doi.org/10.5747/ca.2017.v13.n2.a157>

SALES, L. **Ação de produtos stoller sobre a qualidade de grãos de soja**. 2017. Monografia (Graduação em Agronomia) – Universidade Federal de Mato Grosso, Campus de Sinop, 2017.

SOUZA, S. R. **Beneficiamento e análise de qualidade de grãos de milho e soja na fazenda Campo Bom.** Monografia (Graduação) - Universidade Federal de Santa Catarina, Centro de Ciências Agrárias, 2012.

TEIXEIRA, G. V. **Avaliação das perdas qualitativas no armazenamento da soja.** Dissertação (Mestrado), Campinas, 2001.

TSUKAHARA, R. Y.; FONSECA, I. C. B.; SILVA, M. A. A.; KOCHINSKI, E.G.; PRESTES NETO, J. E.; SUYAMA, J. T. Produtividade de soja em consequência do atraso da colheita e de condições ambientais. **Pesquisa Agropecuária Brasileira**, v.51, n.8, p.905-915, 2016.
<https://doi.org/10.1590/S0100-204X2016000800002>

VINHOTE, J. A.; CORADINE, G.; SILVA, E. B. R.; SILVA, W. C.; SOARES, G. S. **Aspectos relacionados à produção de soja no Brasil – uma revisão da literatura.** 2021. Cap.9.
https://doi.org/10.36599/editpa-2021_caiis-008

XAVIER, T. S.; DARONCH, D. J.; PELUZIO, J. M.; AFFÉRI, F. S.; CARVALHO, E. V.; SANTOS, W. F. Época de colheita na qualidade de grãos de genótipos de soja. **Comunicata Scientiae**, v.6, p.241-245, 2015.